

10/11/15

HOME WORK - 1

Anil Bettadahalli  
Channakeshava

$$-u'' = f \text{ in } ]0, 1[$$

or

$$u'' + f(x) = 0, ]0, 1[$$

with B.C  $u(0) = 0, u(1) = \alpha$

\* Strong form to weak form : using (WRM)

$$\int_0^1 w_i [u'' + f(x)] dx = 0$$

$$\int_0^1 w_i u'' dx + \int_0^1 w_i f(x) dx = 0$$

Integrating by parts

$$\int_0^1 w_i u'' dx = w_i u' \Big|_0^1 - \int_0^1 \frac{dw_i}{dx} \left( \frac{du}{dx} \right) dx$$

$$\int_0^1 \frac{dw_i}{dx} \frac{du}{dx} dx = \int_0^1 w_i f(x) dx + w_i \left( \frac{du}{dx} \right) \Big|_0^1$$

$R$  &  $\bar{R}$  are unknown



Let  $R = -\frac{du}{dx}$  (Unknown) (Reaction force on Dirichlet Boundary) at  $(x=0)$

$\bar{R} \rightarrow$  Reaction force at Neumann Boundary = unknown at  $(x=1)$

$$\int_0^1 \frac{dw_i}{dx} \frac{du}{dx} dx = \int_0^1 w_i f(x) dx + [w_i R]_0 - [w_i \bar{R}]_1 \left. \vphantom{\int_0^1} \right\} \text{Weak form}$$

Approximation of unknown, we have

$$u \approx u^h = \sum_{j=1}^n N_j(x) u_j$$

$$\int_0^1 \frac{dw_i}{dx} \frac{d}{dx} \left[ \sum_{j=1}^n N_j(x) u_j \right] dx = \int_0^1 w_i f(x) dx + [w_i \bar{R}]_0 - [w_i \bar{R}]_1$$

Using Galerkin Method

$$w_i = N_i$$

$$\int_0^1 \frac{dw_i}{dx} \sum_{j=1}^n \frac{d}{dx} [N_j(x)] u_j dx = \int_0^1 N_i f(x) dx + [N_i \bar{R}]_0 - [N_i \bar{R}]_1$$

$\underbrace{\int_0^1 \frac{dw_i}{dx} \sum_{j=1}^n \frac{d}{dx} [N_j(x)] u_j dx}_{K_{ij}} = \underbrace{\int_0^1 N_i f(x) dx + [N_i \bar{R}]_0 - [N_i \bar{R}]_1}_{f_i}$



$$\underline{K} \underline{a} = \underline{f}$$

$$K_{ij} = \frac{dN_i}{dx} \frac{d}{dx} [N_j(x)]$$


$$a = u_j$$

$$f_i = N_i f(x) + [N_i \bar{R}]_0 - [N_i \bar{R}]_1$$


⊗ Linear System of Equations to be solved




# Shape functions




	<u>Global</u>	<u>Local</u>	
$N_1 = N_1^{(1)}$	}	$0 \leq x \leq 1/3$	
$N_1 = 0$			$1/3 \leq x \leq 2/3$
$N_1 = 0$			$2/3 \leq x \leq 1$



$N_2 = N_2^{(1)}$	}	$0 \leq x \leq 1/3$	
$N_2 = N_1^{(2)}$			$1/3 \leq x \leq 2/3$
$N_2 = 0$			$2/3 \leq x \leq 1$



$N_3 = 0$	}	$0 \leq x \leq 1/3$	
$N_3 = N_2^{(2)}$			$1/3 \leq x \leq 2/3$
$N_3 = N_1^{(3)}$			$2/3 \leq x \leq 1$



$N_4 = 0$	}	$0 \leq x \leq 1/3$	
$N_4 = 0$			$1/3 \leq x \leq 2/3$
$N_4 = N_2^{(3)}$			$2/3 \leq x \leq 1$

From weak form, we have

$$\int_0^1 \frac{dN_i}{dx} \left[ \frac{dN_1}{dx} u_1 + \frac{dN_2}{dx} u_2 + \frac{dN_3}{dx} u_3 + \frac{dN_4}{dx} u_4 \right] dx = \int_0^1 N_i f(x) dx + [N_i R]_0 - [N_i R]_1$$

Global Solution System

$i = 1, 2, 3, \dots$

Local System: (Global solution written in local shape function)

for  $i=1$

$$\int_0^{1/3} \frac{dN_1^{(1)}}{dx} \left[ \frac{dN_1^{(1)}}{dx} u_1 + \frac{dN_2^{(1)}}{dx} u_2 \right] dx = \int_0^{1/3} \underbrace{N_1^{(1)} f(x)}_{f_1^{(1)}} dx + \overbrace{[N_1 \bar{R}]_0}^{\bar{R}} - \underbrace{[N_1 \bar{R}]_1}_{\bar{R}}$$

for  $i=2$

$$\int_0^{1/3} \frac{dN_2^{(1)}}{dx} \left[ \frac{dN_1^{(1)}}{dx} u_1 + \frac{dN_2^{(1)}}{dx} u_2 \right] dx + \int_{1/3}^{2/3} \frac{dN_1^{(2)}}{dx} \left[ \frac{dN_1^{(2)}}{dx} u_2 + \frac{dN_2^{(2)}}{dx} u_3 \right] dx$$

$$= \int_0^{1/3} \underbrace{N_2^{(1)} f(x)}_{f_2^{(1)}} dx + 0 + \int_{1/3}^{2/3} \underbrace{N_1^{(2)} f(x)}_{f_1^{(2)}} dx + 0$$

for  $i=3$

$$\int_{1/3}^{2/3} \frac{dN_2^{(2)}}{dx} \left[ \frac{dN_1^{(2)}}{dx} u_2 + \frac{dN_2^{(2)}}{dx} u_3 \right] dx + \int_{2/3}^1 \frac{dN_1^{(3)}}{dx} \left[ \frac{dN_1^{(3)}}{dx} u_3 + \frac{dN_2^{(3)}}{dx} u_4 \right] dx$$

$$= \int_{1/3}^{2/3} \underbrace{N_2^{(2)} f(x)}_{f_2^{(2)}} dx + \int_{2/3}^1 \underbrace{N_1^{(3)} f(x)}_{f_1^{(3)}} dx + 0$$

for  $i=4$

$$\int_{2/3}^1 \frac{dN_2^{(3)}}{dx} \left[ \frac{dN_1^{(3)}}{dx} u_3 + \frac{dN_2^{(3)}}{dx} u_4 \right] dx = \int_{2/3}^1 \underbrace{N_2^{(3)} f(x)}_{f_2^{(3)}} dx + 0 - \overbrace{[N_3 \bar{R}]_1}^{\bar{R}}$$

The above expression can be written in matrix form as,

$$\begin{bmatrix} K_{11}^{(1)} & K_{12}^{(1)} & 0 & 0 & 0 \\ K_{21}^{(1)} & K_{22}^{(1)} + K_{11}^{(2)} & K_{12}^{(2)} & 0 & 0 \\ 0 & K_{21}^{(2)} & K_{22}^{(2)} + K_{11}^{(3)} & K_{12}^{(3)} & 0 \\ 0 & 0 & K_{21}^{(3)} & K_{22}^{(3)} & 0 \end{bmatrix} \begin{bmatrix} u_1 = 0 \\ u_2 \\ u_3 \\ u_4 = \alpha \end{bmatrix} = \begin{bmatrix} f_1^{(1)} + \bar{R} \\ f_2^{(1)} + f_1^{(2)} \\ f_2^{(2)} + f_1^{(3)} \\ f_2^{(3)} - \bar{R} \end{bmatrix}$$

(i)

∴ Stiffness matrix for 'n' noded element

$$K = \begin{bmatrix} K_{11}^{(1)} & K_{12}^{(1)} & & & & & & & & & \\ K_{21}^{(1)} & K_{22}^{(1)} + K_{11}^{(2)} & K_{12}^{(2)} & & & & & & & & \\ & K_{21}^{(2)} & K_{22}^{(2)} + K_{11}^{(3)} & K_{12}^{(3)} & & & & & & & \\ & & K_{21}^{(3)} & K_{22}^{(3)} + K_{11}^{(4)} & K_{12}^{(4)} & & & & & & \\ & & & \vdots & \vdots & & & & & & \\ & & & & K_{22}^{(n-1)} + K_{11}^{(n)} & K_{12}^{(n)} & & & & & \\ & & & & K_{21}^{(n)} & K_{22}^{(n-1)} & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \end{bmatrix}_{n \times n}$$

Global External force Vector

$$\underline{f} = \begin{bmatrix} f_1^{(1)} + R \\ f_2^{(1)} + f_1^{(2)} \\ f_2^{(2)} + f_1^{(3)} \\ \vdots \\ f_2^{(n-2)} + f_1^{(n-1)} \\ f_2^{(n-1)} - \bar{R} \end{bmatrix}_{n \times 1}, \quad a = \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ \vdots \\ u_{n-1} \\ u_n \end{bmatrix}_{n \times 1}$$

$$u = u^h = N_1 u_1 + N_2 u_2 + N_3 u_3 + N_4 u_4$$

Now, we have


$$K_{ij} = \int_{x^{(1)}} \frac{dN_i^{(1)}}{dx} \frac{dN_j^{(1)}}{dx} dx, \quad f_i = \int_{x^{(1)}} N_i^{(1)} f(x) dx$$

We know that :

$$u^h(x) = a_0 + a_1(x) \quad [\text{polynomial approximation}]$$

③ FE Approximation  $u^h$  for  $n=3$ ,  $f(x) = \sin x$ ,  $\alpha=3$

for  $e=1$ , element ①



$$K_{11}^{\textcircled{1}} = \int_{x^{\textcircled{1}}} \left( \frac{-1}{l^e} \right) \cdot \left( \frac{-1}{l^e} \right) dx$$

$$K_{11}^{\textcircled{1}} = \frac{1}{l^e} = K_{22}^{\textcircled{1}}$$

$$K_{12}^{\textcircled{1}} = \int_{x^{\textcircled{1}}} \left( \frac{-1}{l^e} \right) \left( \frac{1}{l^e} \right) dx = \frac{-1}{l^e} = K_{21}^{\textcircled{1}}$$

$$K_{11}^{\textcircled{1}} = K_{22}^{\textcircled{1}} = \int_0^{1/3} \left( \frac{-1}{1/3} \right) \left( \frac{-1}{1/3} \right) dx = \frac{3}{1}$$

$$K_{21}^{\textcircled{1}} = K_{12}^{\textcircled{1}} = \int_0^{1/3} \left( \frac{-1}{1/3} \right) \left( \frac{1}{1/3} \right) dx = -\frac{3}{1}$$

$$f_1^{\textcircled{1}} = \int_0^{1/3} \left( \frac{x_2^{\textcircled{1}} - x}{1/3} \right) \sin x dx = 3 \int_0^{1/3} (x_2^{\textcircled{1}} \sin x - x \sin x) dx$$

$$= 3 \int_0^{1/3} \left[ \frac{1}{3} \sin x - x \sin x \right] dx$$

$$= 3 \left[ -\cos x \Big|_0^{1/3} \right] - 3 \left( -x \cos x + \sin x \right) \Big|_0^{1/3}$$

$$= 0.055043 - 3 \times 0.012209$$

$$\boxed{f_1^{\textcircled{1}} = 0.0184160}$$


$$\begin{aligned}
 f_2^{(1)} &= \int_0^{1/3} \left( \frac{x - x_1^{(1)}}{1/3} \right) \sin x \, dx \\
 &= 3 \int_0^{1/3} [x \sin x - x_1^{(1)} \sin x] \, dx \quad (x_1^{(1)} = 0) \\
 &= 3 \int_0^{1/3} (x \sin x - 0) \, dx \\
 &= 3 \times 0.012209
 \end{aligned}$$

$$\boxed{f_2^{(1)} = 0.036627}$$

\* For  $e = 2$ , Element ②

$$K_{11}^{(2)} = K_{22}^{(2)} = 3$$

$$K_{12}^{(2)} = K_{21}^{(2)} = -3$$



$$\begin{aligned}
 f_1^{(2)} &= \int_{1/3}^{2/3} \left( \frac{x_2^{(2)} - x}{1/3} \right) \sin x \, dx \\
 &= 3 \int_{1/3}^{2/3} (x_2^{(2)} \sin x - x \sin x) \, dx \\
 &= 3 \int_{1/3}^{2/3} \left[ \frac{2}{3} \sin x - x \sin x \right] \, dx \\
 &= -\cos x \Big|_{1/3}^{2/3} - 3 \left[ -x \cos x + \sin x \right] \Big|_{1/3}^{2/3} \\
 &= 2 [0.15907] + 3 [0.082236]
 \end{aligned}$$

$$\boxed{f_1^{(2)} = 0.071432}$$

$$f_2^{(2)} = \int_{1/3}^{2/3} \left( \frac{x - x_1^{(2)}}{1/3} \right) \sin x \, dx$$


$$= 3 \int_{1/3}^{2/3} \left[ x \sin x - \frac{1}{3} \sin x \right] dx$$

$$= 3 \left[ \left[ -x \cos x + \sin x \right] \Big|_{1/3}^{2/3} + \frac{1}{3} \cos x \Big|_{1/3}^{2/3} \right]$$

$$= 3(0.082236) - 0.15907$$

$$\boxed{f_2^{(2)} = 0.087638}$$

\* For  $e=3$ , Element (3).



$$f_1^{(3)} = \int_{2/3}^1 \left( \frac{x_2^{(3)} - x}{1/3} \right) \sin x \, dx$$

$$= 3 \int_{2/3}^1 (x_2^{(3)} \sin x - x \sin x) \, dx$$

$$= 3 \int_{2/3}^1 (\sin x - x \sin x) \, dx$$

$$= 3 \int_{2/3}^1 \sin x \, dx - 3 \int_{2/3}^1 x \sin x \, dx$$

$$= 3 (-\cos x) \Big|_{2/3}^1 - 3 (-x \cos x + \sin x) \Big|_{2/3}^1$$

$$= (3 \times 0.24558) - 3(0.20672)$$

$$\boxed{f_1^{(3)} = 0.116594}$$



$$\begin{aligned}
 f_2^{(3)} &= \int_{2/3}^1 \left( \frac{x - x_1^{(3)}}{1/3} \right) \sin x \, dx \\
 &= 3 \int_{2/3}^1 x \sin x \, dx - 3 \int_{2/3}^1 \frac{2}{3} \sin x \, dx \\
 &= 3 \times [-x \cos x + \sin x] \Big|_{2/3}^1 - 2 [-\cos x] \Big|_{2/3}^1 \\
 &= 3 \times 0.20672 - 2(0.2455)
 \end{aligned}$$

$$\boxed{f_2^{(3)} = 0.12899}$$

Substituting all these values in Eqn (i)

$$3 \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{bmatrix} = \begin{bmatrix} 0.0184160 + R \\ 0.036627 + 0.071432 \\ 0.087638 + 0.116594 \\ 0.12899 - \bar{R} \end{bmatrix}$$

From data,  $u(0) = u_1 = 0$

$u(1) = u_4 = \alpha = 3$

Now, we have

$$3 \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ u_2 \\ u_3 \\ 3 \end{bmatrix} = \begin{bmatrix} 0.0184160 + R \\ 0.036627 + 0.071432 \\ 0.087638 + 0.116594 \\ 0.12899 - \bar{R} \end{bmatrix}$$

$(0.108059)$   
 $(0.204232)$

To solve for  $u_2$  &  $u_3$

$$0 + 6u_2 - 3u_3 = 0.108059$$

$$(0 - 3u_2 + 6u_3 - 9 = 0.204232) \times 2$$

$$\begin{array}{r} 6u_2 - 3u_3 = 0.108059 \\ (+) \quad -6u_2 + 12u_3 = \text{~~0.204232~~} \\ \hline \phantom{6u_2} - 9u_3 = 18.40846 \end{array}$$

$$9u_3 = 18.5165246$$

$$\Rightarrow \boxed{u_3 = 2.0573}$$

$$6u_2 - 3(2.0573) = 0.108059$$

$$\boxed{u_2 = 1.0467}$$

$$\therefore \begin{array}{l} u_1 = 0 \\ u_2 = 1.0467 \\ u_3 = 2.0573 \\ u_4 = 3 \end{array}$$

→ FEM Solution

~~Find~~ ~~Find~~, Find unknowns :  $R$  &  $\bar{R}$  [Reaction forces]

$$3(u_1 + u_2) = 0.0184160 + R$$

$$-3u_2 = 0.0184160 + R$$

$$u_1 = 0$$

$$-3(1.0467) = 0.0184160 + R$$

$$\Rightarrow \boxed{R = -3.1585}$$

$$3(-u_3 + u_4) = 0.12899 - \bar{R}$$

$$3(-2.0573 + 3) = 0.12899 - \bar{R}$$

$$\Rightarrow \boxed{\bar{R} = -2.6991}$$

\* From Data, Given Exact solution is :-

$$u(x) = \sin x + (3 - \sin 1)x$$

→ put  $x = 0$

$$u(0) = \sin 0 + (3 - \sin 1) \cdot 0$$

$$\boxed{u(0) = 0}$$

→ put  $x = \frac{1}{3}$

$$\begin{aligned} u\left(\frac{1}{3}\right) &= \sin\left(\frac{1}{3}\right) + (3 - \sin 1) \cdot \frac{1}{3} \\ &= 0.327194 + (3 - 0.8414) \cdot \frac{1}{3} \end{aligned}$$

$$\boxed{u\left(\frac{1}{3}\right) = 1.04670}$$

→ put  $x = \frac{2}{3}$

$$\begin{aligned} u\left(\frac{2}{3}\right) &= \sin\left(\frac{2}{3}\right) + (3 - \sin 1) \cdot \frac{2}{3} \\ &= 0.61836 + (3 - 0.8414) \cdot \frac{2}{3} \end{aligned}$$

$$\boxed{u\left(\frac{2}{3}\right) = 2.05738}$$

→ put  $x = 1$

$$\begin{aligned} u(1) &= \sin(1) + (3 - \sin 1) \cdot 1 \\ &= 0.8414 + (3 - 0.8414) \end{aligned}$$

$$\boxed{u(1) = 3}$$

∴

$$\begin{aligned} u(0) &= 0 \\ u\left(\frac{1}{3}\right) &= 1.04670 \\ u\left(\frac{2}{3}\right) &= 2.05738 \\ u(1) &= 3 \end{aligned}$$

→ Exact Solution

\* Comparison :-

FEM Solution	Exact Solution	Relative Error
$u_1 = 0$	$u(0) = 0$	0
$u_2 = 1.0467$	$u\left(\frac{1}{3}\right) = 1.0467$	0
$u_3 = 2.0573$	$u\left(\frac{2}{3}\right) = 2.05738$	$8 \times 10^{-5}$
$u_4 = 3$	$u(1) = 3$	0

## COMPARISON BETWEEN FEM SOLUTION AND EXACT SOLUTION

